

## Towards a Catalyst for Alkene Hydroamination - Static and Dynamic Ab Initio DFT Studies

H.M. Senn, P.E. Blöchl, A. Togni

published in

*Modern Methods and Algorithms of Quantum Chemistry*,  
J. Grotendorst (Ed.), John von Neumann Institute for Computing,  
Jülich, NIC Series, Vol. 2, ISBN 3-00-005746-3, p. 45, 2000.

© 2000 by John von Neumann Institute for Computing

Permission to make digital or hard copies of portions of this work for personal or classroom use is granted provided that the copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. To copy otherwise requires prior specific permission by the publisher mentioned above.

<http://www.fz-juelich.de/nic-series/>

# Towards a Catalyst for Alkene Hydroamination – Static and Dynamic *Ab Initio* DFT Studies

Hans Martin Senn,<sup>\*a</sup> Peter E. Blöchl,<sup>b</sup> Antonio Togni<sup>a</sup>

<sup>a</sup> Laboratory of Inorganic Chemistry, ETH Zürich, CH-8092 Zürich, Switzerland


<sup>b</sup> Zürich Research Laboratory, IBM Research Division, CH-8803 Rüschlikon, Switzerland

Using density functional theory, we have investigated group 9 and group 10 d<sup>8</sup> transition-metal complexes in view of their potential application as catalysts for the **hydroamination of alkenes** [1]. We used the **projector-augmented wave (PAW) method** [2], a plane-waves-based method with all-electron wavefunctions capable of performing **Car–Parrinello *ab initio* molecular dynamics** simulations.

Transition states were located using friction dynamics combined with a moving constraint to drive the system over the barrier, while all other degrees of freedom were being relaxed. Dynamical reaction paths were obtained by letting the system evolve freely from the transition state, thus allowing a detailed analysis of the time evolution of the reaction event [3].

The principal mechanistic pathway explored involves external nucleophilic attack of the amine on the coordinated alkene (C=C activation), followed by either protonolytic cleavage of the metal–alkyl bond or protonation of the metal with subsequent C–H reductive elimination. Complexes of the type {MCl(PH<sub>3</sub>)<sub>2</sub>}<sup>n+</sup> have been studied, where M = Ni(II), Pd(II), Pt(II) (*n* = 1) or M = Co(I), Rh(I), Ir(I) (*n* = 0).

The reaction profiles for the complete catalytic cycle will be discussed, assessing the suitability of the metal complexes. The influence of ligand electronic effects on the rate-determining step has been investigated by replacing PH<sub>3</sub> by a series of substituted phosphanes PR<sub>3</sub> (R = F, CF<sub>3</sub>, Me, NMe<sub>2</sub>, Ph, *p*-C<sub>6</sub>H<sub>4</sub>OMe, *p*-C<sub>6</sub>H<sub>4</sub>NMe<sub>2</sub>) with different electron-donating capabilities.



[1] For a recent review, see: T. E. Müller, M. Beller, *Chem. Rev.* **1998**, 98, 675–703.

[2] P. E. Blöchl, *Phys. Rev. B* **1994**, 50, 17953–17979.

[3] P. E. Blöchl, H. M. Senn, A. Togni, in *Transition State Modeling for Catalysis*, D. G. Truhlar, K. Morokuma, Eds., ACS Symposium Series 721, American Chemical Society, Washington, DC, 1999, pp. 88–99.